arXiv:1203.0472v1 [cond-mat.mtrl-sci] 2 Mar 2012

On hereditary models of polymers

Monica De Angelis

Università Studi di Napoli Federico II, Facoltà di Ingegneria,
Dip. Mat. e Appl., via Claudio 21, 80125, Napoli; Italy.
Tel. 081 7683387 e-mail modeange@unina.it

Abstract

An equivalence between an integro-differential operator $\mathcal{M}$ and an evolution operator $\mathcal{L}_n$ is determined. Owing to this equivalence, the fundamental solution of $\mathcal{L}_n$ is estimated in terms of the fundamental solution related to the third order operator $\mathcal{L}_1$ whose behaviour is now acquired. Moreover, properties typical of wave hierarchies can be applied to polymeric materials. As an example the case $n = 2$ is considered and results are applied to *the Rouse model* and *the reptation model* which describe different aspects of polymer chains.

1. Introduction

The creep and relaxation processes related to the viscoelastic behavior of many actual polymeric materials are specified by means of memory functions like:

$$(1.1) \qquad g_n(t) = \sum_{h=1}^{n} B_h \, e^{-\beta_h t},$$

where n, B_h and β_h depend on the polymer physics and are determined in order to fit the experimental curves for $g_n(t)$ with a required approximation [1-3].

Moreover, according to the well-known Muntz and Schwartz'z theorem [4-5], every continuous spectrum can be uniformly represented by Dirichlet polynomials.

In this paper, an integro - differential operator $\mathcal{M}$ related to (1.1) is considered and a *conditioned* equivalence between $\mathcal{M}$ and the $(2+n)$ order operator:

$$(1.2) \qquad \mathcal{L}_n = \partial_t^{(n)}(\partial_{tt} - c_n^2 \partial_{xx}) +$$

$$a_{n-1}\partial_t^{(n-1)}(\partial_{tt} - c_{n-1}^2 \partial_{xx}) + ..a_0 \, (\partial_{tt} - c_0^2 \partial_{xx})$$

where a_k $(k = 0..n-1)$ are positive constants, is extablished.

By this equivalence whatever n may be the fundamental solution of (1.2) is explictly determined and an estimate in terms of the fundamental solution related to the third order operator is achieved, too.

As applications, when $n = 2$, results are applied to polymer chains and *the Rouse model* and *the reptation model* are considered.

2. Statement of the problem

Let $\mathcal{B}$ a linear, isotropic, homogeneous system and let $\underline{u}(x,t)\underline{i}$ the displacement field from an underformed reference configuration $\mathcal{B}_0$.

Indicating by σ and ε the only non vanishing components of the stress and the strain tensors, the constitutive relation is

$$\sum_{k=0}^{n} a_k \ \partial_t^k \sigma = \sum_{k=0}^{n} \alpha_k \ \partial_t^k \varepsilon \tag{2.1}$$

with a_k, α_k constant $(a_n, \alpha_n \neq 0)$.
So, if $\underline{\mathbf{f}} = f\underline{\mathbf{i}}$ is the known body force, ρ_0 denotes the mass density in $\mathcal{B}_0$, and $c_k = \alpha_k/\rho_0 a_k$, the one dimensional linear motions of $\mathcal{B}$ are described by higher order equation [6]:

$$\mathcal{L}_n u = F \tag{2.2}$$

where

$$\mathcal{L}_n = \sum_{k=0}^{n} a_k \partial_t^{(k)} (\partial_{tt} - c_k \partial_{xx}) \tag{2.3}$$

and

$$F = (1/\rho_0) \ \sum_{k=0}^{n} a_k \ \partial_t^k f. \tag{2.4}$$

In (2.3) constants c_k are the characterized speeds depending on the materials properties of the medium and in many physical problems it results $c_0^2 < c_1^2 .. < c_{n-1}^2 < c_n^2$ and so the equation (1.2) is typical of *the wave hierarchies.* [7].

When $n = 1$, (2.3) turns into a strictly hyperbolic third - order operator which models the evolution of the Standard Linear Solid (S.L.S.) [8-9] and its behaviour has been already discussed in [10]. The fundamental solution $\mathcal{E}_1$ has been explicitly determined together with maximum theorems and boundary layer estimates.

Moreover, if $J(t)$ denotes the creep-compliance, the behavior of most viscoelastic media is fairly well modelled by linear hereditary equations like:

$$\varepsilon(t) \ = \ J(0) \quad [\sigma(t) + \\ + \int_{-\infty}^{t} \dot{J}(t-\tau)\sigma(\tau)d\tau]. \tag{2.5}$$

According to *fading memory* hypotheses [11-12], $\dot{J}(t)$ is a positive fast decreasing function and in many real materials as polymers rubbers or bitumines, it is representated by means of chains of S.L.S. elements in series or parallel [2-3]. In the series case, one has:

$$J_n(t) = \ J_n(0) \ \ [\ 1+ \tag{2.6}$$

$$+ \sum_{k=1}^{n} \frac{B_k}{\beta_k}(1 - e^{-\beta_k t})],$$

where n is the number of elements in the chain, $J_n(0)$ denotes the elastic compliances and $\tau_k = \beta_k^{-1}$ are the characteristic times .

Let consider equations of one dimensional motions of $\mathcal{B}$:

$$\rho_0 \, u_{tt} = \sigma + f, \quad \varepsilon = u, \tag{2.7}$$

by means of (2.5) - (2.7) the following integral constitutive equation is deduced:

$$\mathcal{M}u = c^2 u_{xx} - u_{tt} - \tag{2.8}$$

$$- \int_0^t g(t-\tau) u_{\tau\tau} d\tau = -F_*(x,t)$$

where

$$c^2 = [\rho_0 J_n(0)]^{-1}, \tag{2.9}$$

$$F_* = c^2 [J_n(0) f + \int_{-\infty}^{0} \dot{J}_n(t-\tau) \sigma_x(\tau) d\tau].$$

and

$$g \; = \; g_n(t) = \sum_{k=1}^{n} B_k e^{-\beta_k t} \; = \dot{J}_n(t)/J_n(0). \tag{2.10}$$

In this memory function, n is quite arbitrary and constants B_k and frequencies β_k are such that:

$$0 < \beta_1 < \beta_2 ... < \beta_n; \tag{2.11}$$

$$B_k > 0 \;\; \forall k = 1, 2....n.$$

Whatever n may be, the fundamental solution E_n of operator $\mathcal{M}$ has been explictly determined [13-14]. Moreover, let E_1 the fundamental solution related to an appropriate S.L.S. $\mathcal{B}_1^*$ defined by:

$$g_1 = b \; e^{-\beta_1 t} \tag{2.12}$$

with

$$b = \beta_1 \sum_1^n \frac{B_k}{\beta_k}, \tag{2.13}$$

the following theorem assures that the fundamental solution E_n can be rigorously estimated by means of E_1.

In fact, if Γ is the open forward characteristic cone $\{(t,x) \, : \, t > 0 \; |x| < ct\}$, and $\chi_n = \prod_{k=2}^{n} (\frac{B_k}{\beta_1})^2$, then the following theorem holds:

Theorem 1.1 - *If the memory function is given by* (2.10) (2.11), *then the fundamental solution* E_n *of* $\mathcal{M}$ *is a never negative* $C^\infty(\Gamma)$ *function and it satisfies the estimate*:

$$0 < E_n(\beta_1..\beta_n, B_1..B_n) < \tag{2.14}$$

$$< \chi_n \;\; E_1(\beta_1, b),$$

everywhere in the cone Γ *and whatever* n *may be.* ∎

3 . Conditioned equivalence between operators $\mathcal{L}_n$ and $\mathcal{M}$

Let be null the initial data related to (2.2) and (2.8) and let

$$\mu_k = a_k/a_n, \tag{3.1}$$

$$\lambda_k = a_k c_k/a_n c_n \quad (k = 0, ..n).$$

Appling the Laplace transform and the polinomial identity one has $c_n = c^2$ and:

$$\left\{ \begin{array}{l} \lambda_0 = \beta_1\beta_2..\beta_n \\ .... \\ \lambda_{n-2} = \beta_1\beta_2 + \beta_1\beta_3 + ..\beta_{n-1}\beta_n \\ \\ \lambda_{n-1} = \beta_1 + ... + \beta_n \end{array} \right. \tag{3.2}$$

So, owing to (2.11), all the λ_k's are positive. Further, as for μ_k, one has:

$$\left\{ \begin{array}{l} \mu_0 = \lambda_0 + B_1(\beta_2..\beta_n) + ... \\ \qquad ... + B_n(\beta_1..\beta_{n-1}) \\ .... \\ \mu_{n-2} = \lambda_{n-2} + B_1(\beta_2 + .. + \beta_n) \\ \qquad ... + B_n(\beta_1 + .. + \beta_{n-1}) \\ \\ \mu_{n-1} = \lambda_{n-1} + B_1 + ... + B_n \end{array} \right. \tag{3.3}$$

and (2.11), (3.3) imply too:

$$0 < \lambda_k < \mu_k \quad (k = 0, ..n - 1). \tag{3.4}$$

So, for $k = 0, ...n - 1$, one has:

$$0 < c_k < c_n = c^2. \tag{3.5}$$

At last, by (3.2),(3.3), it follows:

$$\frac{\lambda_0}{\mu_0} < \frac{\lambda_1}{\mu_1} < \frac{\lambda_{n-1}}{\mu n - 1} \tag{3.6}$$

and hence:

$$0 < c_0 < c_1.... < c_n. \tag{3.7}$$

As consequence, the following property holds:

Property 3.1 *Hypotheses of fading memory* (2.10) (2.11) *imply that the differential operator* (2.3) *is typical of wave hierarchies.* ■

Vice versa when the differential equation (2.2) is prefixed, to obtain the dual hereditary equation (2.8) with a memory function $g_n(t)$ satisfying (2.10), (2.11), appropriate restrictiones on the constants a_k, c_k must be imposed.

Example 3.1

When $n = 2$, one has $c^2 = c_2, B_0 = 1$. Further β_1, β_2 are real iff:

$$\omega^2 = (a_1c_1)^2 - 4(a_0c_0)(a_2c_2) > 0. \tag{3.8}$$

So, beeing:

$$\beta_1 = \frac{1}{2a_2c_2}(a_1c_1 - \omega), \tag{3.9}$$

$$\beta_2 = \frac{1}{2a_2c_2}(a_1c_1 + \omega),$$

it results $0 < \beta_1 < \beta_2$.
Moreover as for $B_i (i = 1, 2)$, one has:

$$(3.10) \quad B_i = \frac{(-1)^{i-1}}{\omega} [a_0(c_2 - c_0) -$$

$$-a_1 \beta_i (c_2 - c_1)], \quad (i = 1, 2),$$

and hence, $B_1 > 0, B_2 > 0$ iff

$$(3.11) \qquad \beta_1 < \frac{a_o}{a_1} \ \frac{c_2 - c_0}{c_2 - c_1} < \beta_2.$$

Therefore, the fourth-order operator:

$$(3.12) \qquad a_2(u_{tt} - c_2 u_{xx})_{tt} \ +$$

$$+ a_1(u_{tt} - c_1 u_{xx})_t + a_0(u_{tt} - c_0 u_{xx})$$

can be analyzed by integral operator $\mathcal{M}$ (2.8) with conditions (2.10), (2.11) and when the constants $a_k, \ c_k$ satisfy (3.8) and (3.11). ∎

4 . Polymeric materials

Polymeric materials are very flexible like rubber and are easily formed into fibres, thin films, etc. Moreover, the liquid state composed only of polymers (polymer melt) is an important state for industrial uses where polymeric materials are processed into various plastic products such as gaskets, seals, flexible joints, vehicle tires, etc.. Also the durability is a requirement imposed on polymers and polymeric composites and the interest for future development of these materials is increasing more and more.

A large literature treats with polymer physics and as for viscoelastic theories, two models which describe different aspects of the polymer chains, have met with reasonable success: *the Rouse model* and *the reptation model* [2-3].

In both cases the memory function $g_n(t)$ assumes a form like (1.1).

In fact in *the Rouse model* function $g_n(t)$ is given by:

$$(4.1) \qquad g_n(t) = k_1 \sum_{h=1}^{n} e^{2h^2 \frac{t}{\tau_1}}$$

where the relaxation time τ_1 can be calculated by means of experimental results.[3].

When the viscoelastic behaviour is represented by *the reptation model*, as times increases, the stress function decreases with a relaxation time τ_d, and one has:

$$(4.2) \qquad g_n(t) = k \sum \frac{1}{h^2} \ e^{-h^2 \frac{t}{\tau_d}},$$

where h ranges over odd integer, the constant k depends on the polymer physics and the value of the *reptation* time τ_d can be fixed according to elasticity experiments [1].

So, if one considers the first two steps in *the reptation model*, it results: $B_1 = k, \ \ B_2 = B_1/9, \ \ \beta_1 = 1/\tau_d, \ \ \beta_2 = 9\beta_1$. Consequently the operator (3.12) is characterized by constants:

$$\begin{cases} c_0 = c^2\,\frac{81}{81+82k\tau_d} & a_0 = 1 + \frac{82}{81}\,k\tau_d \\ c_1 = c^2\,\frac{9}{9+k\tau_d} & a_1 = \frac{10\tau_d^2}{9}(\frac{1}{\tau_d} + \frac{k}{9}) \\ c_2 = c^2 & a_2 = \frac{\tau_d^2}{9} \end{cases} \tag{4.3}$$

Analogously, in *the Rouse model*, beeing $B_1 = B_2 = k_1, \;\; \beta_1 = 2/\tau_1, \;\; \beta_2 = 4\beta_1$, one has:

$$\begin{cases} c_0 = c^2\frac{8}{8+5k_1\tau_1} & a_0 = 1 + \frac{5k_1}{8}\tau_1 \\ c_1 = c^2\frac{5}{5+k_1\tau_1} & a_1 = \frac{\tau_1^2}{16}(2k_1 + \frac{10}{\tau_1}) \\ c_2 = c^2 & a_2 = \frac{\tau_1^2}{16} \end{cases} \tag{4.4}$$

The *wave hierarchies* defined by (4.3) or (4.4) are governed by the operator $\mathcal{L}_1^*$ of the Standard Linear Solid defined, respectively, by:

$$\text{(4.5)}\begin{cases} c_0 = c^2\,\frac{81}{81+82k\tau_d} & c_1 = c^2 \\ a_0 = 1 + \frac{82}{81}\,k\tau_d & \eta = \frac{81\tau_d}{81+82k\tau_d}, \end{cases}$$

$$\text{(4.6)}\begin{cases} c_0 = c^2\frac{8}{8+5k_1\tau_1} & c_1 = c^2 \\ a_0 = 1 + \frac{5}{8}k_1\tau_1 & \eta = \frac{4\tau_1}{8+5k_1\tau_1}. \end{cases}$$

Remark 3.1- As swowed, memory function $g_n(t)$ can depend on h^2. So the approximation to the two first terms appears to be reasonable. However, in many article the model is limited to a single relaxtion time (see. f.i.[15]). ∎

References

1. Ferry, J. D.: Viscoelastic properties of polymers. New York. Wiles, 1961
2. Doi, M. See, H.: Introduction to Polymer Physics. Oxford: Clarendon Press, 1977
3. Doi, M. Edwards S. F.: The theory of Polymer Dynamics. Oxford: Clarendon Press, 1986
4. Sunyer Balaguer, F. : Approximation of functions by linear combinations of exponentials. Collect. Math. **17**, 145-177, (1965).
5. Schwartz, L.: Etude des sommes d'exponentielles. Act. Scient. et Industr. **959**, Hermann, Paris (1959).
6. Christensen, R. M.: Theory of Viscolasticity. New York : Academic Press, 1971
7. Whitham, G. B.: Linear and Non linear Waves. New York : John Wiley e Sons, 1974
8. Hunter, S.C.: Mechanics of Continuous media. England: Ellis Horwood, 1976

9. Haupt, P.: Continuous Mechanics and theory of Materials, Springer 2000

10. Renno, P.: On a Wave Theory for the Operator $\varepsilon\partial_t(\partial_t^2 - c_1^2\Delta_n) + \partial_t^2 - c_0^2\Delta_n$. Ann. Mat. pura e Appl.,**136(4)** 355-389 (1984).

11. Graffi. D. : On the fading memory. Applicable Analysis, **15** (1983) 295-311

12. Graffi. D.: Mathematical models and waves in linear viscoestacity. Euromech Colloquium 127 on *Waves propagation in viscoelastic media*, Pitman Adv. Publ. Comp. Research notes in Math.,**52**, 1-27 1980

13. Renno, P.: On the Cauchy problem il linear viscoelasticity. Ren. Acc. Naz. Lincei, VIII vol. **LXXXV** (1983)

14. Renno, P.: On some viscoelastic models. Ren. Acc. Naz. Lincei, VIII vol. **LXXV** (1983)

15. Ianniruberto,G., Marrucci G.: A simple constiyutive equation for entangled polymers with chain stretch. J. Rheol.**45**(6) 1305- 1318 (2001)